

$D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0$ DECAY WIDTH

Marina Nielsen

Instituto de Física, Universidade de São Paulo, C.P. 66318, 05389-970 São Paulo, SP, Brazil

We use the QCD sum rules to analyze the hadronic decay $D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0$, in the hypothesis that the $D_{sJ}^+(2317)$ can be identified as a four-quark state. We use a diquark-antidiquark current and work to the order of m_s in full QCD, without relying on $1/m_c$ expansion. We find that the partial decay width of the hadronic isospin violating mode is proportional to the isovector quark condensate, $\langle 0 | \bar{d}d - \bar{u}u | 0 \rangle$. The estimated partial decay width is of the order of 6 keV.

PACS numbers: 11.55.Hx, 12.38.Lg, 13.25.-k

The strange-charmed mesons $D_{sJ}^+(2317)$ and $D_{sJ}^+(2460)$ with spin parity 0^+ , 1^+ [1, 2, 3, 4] are lighter than the prediction of the very successful quark model for the charmed mesons [5]. One interpretation is that this is evidence for a chiral symmetry where the mass gap between the 0^- and 1^- states equates the mass gap between the 0^+ and 1^+ states [6]. Other interpretations are related with the underlying structure of these mesons, which has been extensively debated. They have been interpreted as conventional $c\bar{s}$ states [7, 8, 9, 10, 11, 12, 13, 14, 15, 16], two-meson molecular state [17, 18], $D - K$ - mixing [19] or four-quark states [20, 21, 22, 23, 24, 25, 26, 27].

Because of their low masses, these two states are lower than the DK and D^*K thresholds. Therefore, their strong decays must proceed through isospin violating effects. There have been some discussions of their decays within the quark model [6, 13, 14, 15, 20] and QCD sum rules [16]. In all these studies but [14], the isospin violating effects were considered through the $\eta - \pi^0$ mixing. However, if these mesons are considered as four-quark states, in a QCD sum rule calculation, the isospin violating effects can be introduced through the mass and quark condensate difference between the u and d quarks.

In a recent calculation [23] the scalar-isoscalar meson $D_{sJ}^+(2317)$ were considered as a S -wave bound state of a diquark-antidiquark pair. As suggested in ref. [28], the diquark was taken to be a spin zero colour anti-triplet. The corresponding interpolating field is:

$$j_S = \frac{\epsilon_{abc}\epsilon_{dec}}{\sqrt{2}} [(u_a^T C \gamma_5 c_b)(\bar{u}_d \gamma_5 C \bar{s}_e^T) + (d_a^T C \gamma_5 c_b)(\bar{d}_d \gamma_5 C \bar{s}_e^T)], \quad (1)$$

where $a, b, c, \dots$ are colour indices and C is the charge conjugation matrix. In ref. [23], using the QCD sum rule (QCDSR) formalism [29, 30, 31], it was shown that it is possible to reproduce the experimental mass of the meson $D_{sJ}^+(2317)$ using this four-quark state picture. Here, we extend the calculation done in ref. [23] to study the vertex associated with the decay $D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0$.

The QCDSR calculation for the vertex, $D_{sJ}^+(2317) D_s^+ \pi^0$, centers around the three-point function given by

$$\Gamma_\mu(p, p', q) = \int d^4x d^4y e^{ip' \cdot x} e^{iq \cdot y} \langle 0 | T[j_{D_s}(x) j_{5\mu}^{\pi^0}(y) j_S^\dagger(0)] | 0 \rangle, \quad (2)$$

where $p = p' + q$ and the interpolating fields for the pion and D_s mesons are given by:

$$j_{5\mu}^{\pi^0} = \frac{1}{\sqrt{2}} (\bar{u}_a \gamma_\mu \gamma_5 u_a - \bar{d}_a \gamma_\mu \gamma_5 d_a), \quad j_{D_s} = i \bar{s}_a \gamma_5 c_a. \quad (3)$$

The fundamental assumption of the QCD sum rule approach is the principle of duality. Specifically, we assume that there is an interval over which the above vertex function may be equivalently described at both, the quark level and at the hadron level. Therefore, the underlying procedure of the QCD sum rule technique is the following: on one hand we calculate the vertex function at the quark level in terms of quark and gluon fields. On the other hand, the vertex function is calculated at the hadronic level introducing hadron characteristics such as masses and coupling constants. At the quark level the complex structure of the QCD vacuum leads us to employ the Wilson's operator product expansion (OPE). The calculation of the phenomenological side proceeds by inserting intermediate states for D_s , π^0 and D_{sJ} , and by using the definitions:

$$\langle 0 | j_{5\mu}^{\pi^0} | \pi^0(q) \rangle = i q_\mu F_\pi, \quad \langle 0 | j_{D_s} | D_s(p') \rangle = \frac{m_{D_s}^2 f_{D_s}}{m_c + m_s}, \quad \langle 0 | j_{D_{sJ}} | D_{sJ}(p) \rangle = \lambda_S. \quad (4)$$

We obtain the following relation:

$$\Gamma_\mu^{phen}(p, p', q) = \frac{\lambda_S m_{D_s}^2 f_{D_s} F_\pi g_{D_{sJ} D_s \pi}}{(m_c + m_s)(p^2 - m_{D_{sJ}}^2)(p'^2 - m_{D_s}^2)(q^2 - m_\pi^2)} q_\mu + \text{continuum contribution}, \quad (5)$$

where the coupling constant $g_{D_{sJ}D_s\pi}$ is defined by the on-mass-shell matrix element

$$\langle D_s\pi | D_{sJ} \rangle = g_{D_{sJ}D_s\pi}. \quad (6)$$

The continuum contribution in Eq.(5) contains the contributions of all possible excited states.

For the light scalar mesons, considered as diquark-antidiquark states, the study of their vertices functions using the QCD sum rule approach at the pion pole [11, 30, 32, 33], was done in ref.[34]. In Table I we show the results obtained for the different vertices studied in ref. [34], as well as the experimental values.

Table I: Numerical results for the coupling constants

vertex	$g(\text{GeV})$	$g^{exp}(\text{GeV})$
$\sigma\pi^+\pi^-$	3.1 ± 0.5	2.6 ± 0.2
$\kappa K^+\pi^-$	3.6 ± 0.3	4.5 ± 0.4
$f_0 K^+ K^-$	1.6 ± 0.1	
$f_0\pi^+\pi^-$	0.47 ± 0.05	1.6 ± 0.8

From Table I we see that, although not exactly in between the experimental error bars, the hadronic couplings determined from the QCD sum rule calculation are consistent with existing experimental data. The biggest discrepancy is for $g_{f_0\pi^+\pi^-}$ and this can be understood since the $f_0 \rightarrow \pi^+\pi^-$ decay is mediated by gluon exchange and, therefore, probably in this case α_s corrections, which were not considered, could play an important role.

Here, we follow ref. [34] and work at the pion pole. The main reason for working at the pion pole is that the matrix element in Eq.(6) defines the coupling constant only at the pion pole. For $q^2 \neq 0$ one would have to replace $g_{D_{sJ}D_s\pi}$, in Eq.(6), by the form factor $g_{D_{sJ}D_s\pi}(q^2)$ and, therefore, one would have to deal with the complications associated with the extrapolation of the form factor [35, 36]. The pion pole method consists in neglecting the pion mass in the denominator of Eq. (5) and working at $q^2 = 0$. In the OPE side one singles out the leading terms in the operator product expansion of Eq.(2) that match the $1/q^2$ term. In the phenomenological side, in the structure $\frac{q_\mu}{q^2}$ we get:

$$\Gamma^{phen}(p^2, p'^2) = \frac{\lambda_S m_{D_s}^2 f_{D_s} F_\pi g_{D_{sJ}D_s\pi}}{(m_c + m_s)(p^2 - m_{D_{sJ}}^2)(p'^2 - m_{D_s}^2)} + \int_{m_c^2}^{\infty} \frac{\rho_{cont}(p^2, u)}{u - p'^2} du. \quad (7)$$

In Eq.(7), $\rho_{cont}(p^2, u)$, gives the continuum contributions, which can be parametrized as $\rho_{cont}(p^2, u) = \frac{b(u)}{s_0 - p^2} \Theta(u - u_0)$ [37, 38], with s_0 and u_0 being the continuum thresholds for D_{sJ} and D_s respectively. Since we are working at $q^2 = 0$, we take the limit $p^2 = p'^2$ and we apply the Borel transformation to $p^2 \rightarrow M^2$ to get:

$$\Gamma^{phen}(M^2) = \frac{\lambda_S m_{D_s}^2 f_{D_s} F_\pi g_{D_{sJ}D_s\pi}}{(m_c + m_s)(m_{D_{sJ}}^2 - m_{D_s}^2)} \left(e^{-m_{D_s}^2/M^2} - e^{-m_{D_{sJ}}^2/M^2} \right) + A e^{-s_0/M^2} + \int_{u_0}^{\infty} \rho_{cc}(u) e^{-u/M^2} du, \quad (8)$$

where

$$A = - \int_{u_0}^{\infty} \frac{b(u)}{(s_0 - u)} du, \quad \text{and} \quad \rho_{cc}(u) = \frac{b(u)}{(s_0 - u)}, \quad (9)$$

stands for the pole-continuum transitions and pure continuum contributions. For simplicity, one assumes that the pure continuum contribution to the spectral density, $\rho_{cc}(u)$, is given by the result obtained in the OPE side. Asymptotic freedom ensures that equivalence for sufficiently large u . Therefore, one uses the ansatz: $\rho_{cc}(u) = \rho_{OPE}(u)$. In Eq.(8), A is a parameter which, together with $g_{D_{sJ}D_s\pi}$, have to be determined by the sum rule.

In the OPE side we work at leading order and deal with the strange quark as a light one and consider the diagrams up to order m_s . To keep the charm quark mass finite, we use the momentum-space expression for the charm quark propagator. We calculate the light quark part of the correlation function in the coordinate-space, which is then Fourier transformed to the momentum space in D dimensions. The resulting light-quark part is combined with the charm-quark part before it is dimensionally regularized at $D = 4$. Singling out the leading terms proportional to q_μ/q^2 , we can write the Borel transform of the correlation function in the OPE side in terms of a dispersion relation:

$$\Gamma^{OPE}(M^2) = \int_{m_c^2}^{\infty} \rho_{OPE}(u) e^{-u/M^2} du, \quad (10)$$

where the spectral density, ρ_{OPE} , is given by the imaginary part of the correlation function. Transferring the pure continuum contribution to the OPE side, the sum rule for the coupling constant, up to dimension 7, is given by:

$$C \left(e^{-m_{D_s}^2/M^2} - e^{-m_{D_{sJ}}^2/M^2} \right) + A e^{-s_0/M^2} = \gamma \langle \bar{q}q \rangle \left[\frac{1}{24\pi^2} \int_{m_c^2}^{u_0} du e^{-u/M^2} u \left(1 - \frac{m_c^2}{u} \right)^2 \right. \\ \left. + \frac{m_c m_s}{8\pi^2} \int_{m_c^2}^{u_0} du e^{-u/M^2} \left(1 - \frac{m_c^2}{u} \right) - \frac{m_c \mathfrak{B}}{6} e^{-m_c^2/M^2} + \frac{m_s \mathfrak{B}}{12} e^{-m_c^2/M^2} \left(1 + \frac{m_c^2}{M^2} \right) \right], \quad (11)$$

with

$$C = \frac{\lambda_S m_{D_s}^2 f_{D_s} F_\pi}{(m_c + m_s)(m_{D_{sJ}}^2 - m_{D_s}^2)} g_{D_{sJ} D_s \pi}. \quad (12)$$

In Eq.(11), γ measures the isospin symmetry breaking in the quark condensate:

$$\gamma = \frac{\langle 0 | \bar{d}d - \bar{u}u | 0 \rangle}{\langle 0 | \bar{u}u | 0 \rangle}, \quad (13)$$

and is the source of nonperturbative isospin violation in the OPE of the vertex function. The value of γ has been estimated in a variety of approaches [32, 39, 40], with results varying over almost one order of magnitude: $-1 \times 10^{-2} \leq \gamma \leq -2 \times 10^{-3}$. A more recent calculation [41] gives a bigger (in module) value: $\gamma = -2.6 \times 10^{-2}$.

In the numerical analysis of the sum rules, the values used for the meson masses, quark masses and condensates are: $m_{D_{sJ}} = 2.317$ GeV, $m_{D_s} = 1.968$ GeV, $m_s = 0.13$ GeV, $m_c = 1.2$ GeV, $\langle \bar{q}q \rangle = -(0.23)^3$ GeV³, $\langle \bar{s}s \rangle = 0.8 \langle \bar{q}q \rangle$. For the meson decay constants we use $F_\pi = \sqrt{2} 93$ MeV and $f_{D_s} = 0.22$ GeV (obtained using $u_0 = 6$ GeV² [42]). For the current meson coupling, λ_S , defined in Eq.(4) we are going to use the result obtained from the two-point function in ref. [23]. Considering $2.6 \leq \sqrt{s_0} \leq 2.8$ GeV we get $\lambda_s = (2.9 \pm 0.3) \times 10^{-3}$ GeV⁵.

In Fig. 1 we show, through the dots, the right-hand side (RHS) of Eq.(11), for $u_0 = 6$ GeV² and $\gamma = -1 \times 10^{-2}$ [40], as a function of the Borel mass. We use the same Borel window as defined in ref.[23].




FIG. 1: Dots: the RHS of Eq.(11), as a function of the Borel mass. The solid line gives the fit of the QCDSR results through the LHS of Eq.(11).

To determine $g_{D_{sJ} D_s \pi}$ we fit the QCDSR results with the analytical expression in the left-hand side (LHS) of Eq.(11):

$$C \left(e^{-m_{D_s}^2/M^2} - e^{-m_{D_{sJ}}^2/M^2} \right) + A e^{-s_0/M^2}, \quad (14)$$

and we get (using $\sqrt{s_0} = 2.7$ GeV): $C = 8.34 \times 10^{-6}$ GeV⁷ and $A = 6.96 \times 10^{-6}$ GeV⁷. Using the definition of C in Eq.(12) and $\lambda_S = 2.9 \times 10^{-3}$ GeV⁵ (the value obtained for $\sqrt{s_0} = 2.7$ GeV) we get $g_{D_{sJ} D_s \pi} = 50$ MeV. Allowing

s_0 to vary in the interval $2.6 \leq \sqrt{s_0} \leq 2.8$ GeV, the corresponding variation obtained for the coupling constant is $45 \text{ MeV} \leq g_{D_{sJ} D_s \pi} \leq 55 \text{ MeV}$.

Fixing $\sqrt{s_0} = 2.7 \text{ GeV}$ and varying the quark condensate, the charm quark and the strange quark masses in the intervals: $-(0.24)^3 \leq \langle \bar{q}q \rangle \leq -(0.22)^3 \text{ GeV}^3$, $1.1 \leq m_c \leq 1.3 \text{ GeV}$ and $0.11 \leq m_s \leq 0.15 \text{ GeV}$, we get results for the coupling constant still between the lower and upper limits given above. However, varying the value of γ from -1×10^{-2} to the more recent value given in [41]: $\gamma = -2.6 \times 10^{-2}$, and keeping the other parameters fixed we get $g_{D_{sJ} D_s \pi} = 130 \text{ MeV}$. On the other hand, if we use the smallest (in module) value allowed for γ : $\gamma = -2. \times 10^{-3}$, we get $g_{D_{sJ} D_s \pi} = 10 \text{ MeV}$. Therefore, the biggest source of uncertainty in our calculation is the value of γ . In all cases considered here, the quality of the fit between the LHS and the RHS of Eq.(11) is similar to the one shown in Fig.1.

The coupling constant, $g_{D_{sJ} D_s \pi}$, is related with the partial decay width through the relation:

$$\Gamma(D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0) = \frac{1}{16\pi m_{D_{sJ}}^3} g_{D_{sJ} D_s \pi}^2 \sqrt{\lambda(m_{D_{sJ}}^2, m_{D_s}^2, m_\pi^2)}, \quad (15)$$

where $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2ac - 2bc$. Considering $\gamma = -1 \times 10^{-2}$, which was the value found in [40] to be consistent with the neutron-proton mass difference in a QCDSR calculation, and allowing $\langle \bar{q}q \rangle$, m_c , m_s and s_0 to vary in the ranges discussed above we get:

$$\Gamma(D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0) = (6 \pm 2) \text{ keV}. \quad (16)$$

However, it is important to state that, if the value for γ found in ref.[41] proves to be correct, then the partial decay width could be as large as $\Gamma(D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0) \sim 40 \text{ keV}$, in agreement with the QCDSR calculation done in ref.[16], where the meson $D_{sJ}^+(2317)$ is considered as a ordinary $c\bar{s}$ state.

In Table II we show the partial decay width obtained by different theoretical groups. The first five calculations assume a $c\bar{s}$ picture for $D_{sJ}^+(2317)$, while the last two assume a four-quark picture for it.

Table II: The decay width $\Gamma(D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0)$ from various theoretical approaches.

ref.	[6]	[13]	[14]	[15]	[16]	[20]	this work
$\Gamma \text{ (keV)}$	21.5	7 ± 1	~ 10	16	39 ± 5	$10 - 100$	6 ± 2

From the results in Table II we see that we can not get a definitive answer about the structure of the $D_{sJ}(2317)$ meson from its strong decay width, since in both pictures: ordinary $c\bar{s}$ or four-quark states, different approaches can give results varying from a few keV to a hundred keV.

We have presented a QCD sum rule study of the vertex function associated with the strong decay $D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0$, where the $D_{sJ}^+(2317)$ meson was considered as diquark-antidiquark state. We found that the source of isospin violation in our calculation is the parameter $\gamma = \frac{\langle 0 | \bar{d}d - \bar{u}u | 0 \rangle}{\langle 0 | \bar{u}u | 0 \rangle}$, which measures the isospin symmetry breaking in the quark condensate. Since, in our approach, the partial decay width is directly proportional to γ^2 , and since there is a large uncertainty in the value of γ , considering γ in the range $-2.6 \times 10^{-2} \leq \gamma \leq -2 \times 10^{-3}$ we get the partial decay width in the range $0.2 \text{ keV} \leq \Gamma(D_{sJ}^+(2317) \rightarrow D_s^+ \pi^0) \leq 40 \text{ keV}$. However, from other QCDSR calculation, we believe that the value of γ is $\sim -1 \times 10^{-2}$, which gives the result shown in Eq.(16).

As a final remark we would like to point out that if, instead of using a isoscalar current, we have used a isovector current for $D_{sJ}(2317)$ (as suggested in ref. [27]), the difference in Eq.(11) would be a factor 2 in the place of γ . In this case the decay would be isospin allowed and the partial width would be $\sim 230 \text{ MeV}$, much bigger than the experimental upper limit to the total width $\Gamma \sim 5 \text{ MeV}$.

Acknowledgements: I would like to thank F.S. Navarra and I. Bediaga for fruitful discussions. This work has been supported by CNPq and FAPESP.

-
- [1] BABAR Coll., B. Auber *et al.*, Phys. Rev. Lett. **90**, 242001 (2003); Phys. Rev. **D69**, 031101 (2004).
 - [2] CLEO Coll., D. Besson *et al.*, Phys. Rev. **D68**, 032002 (2003).
 - [3] BELLE Coll., P. Krokovny *et al.*, Phys. Rev. Lett. **91**, 262002 (2003).
 - [4] FOCUS Coll., E.W. Vaandering, hep-ex/0406044.

- [5] S. Godfrey and N. Isgur, Phys. Rev. **D32**, 189 (1985); S. Godfrey and R. Kokoshi, Phys. Rev. **D43**, 1679 (1991).
- [6] W. Bardeen, E. Eichten and C. Hill, Phys. Rev. **D68**, 054024 (2003).
- [7] Y.-B. Dai, C.-S. Huang, C. Liu and S.-L. Zhu, Phys. Rev. **D68**, 114011 (2003).
- [8] G.S. Bali, Phys. Rev. **D68**, 071501(R) (2003).
- [9] A. Dougall, R.D. Kenway, C.M. Maynard and C. Mc-Neile, Phys. Lett. **B569**, 41 (2003).
- [10] A. Hayashigaki and K. Terasaki, hep-ph/0411285.
- [11] S. Narison, Phys. Lett. **B605**, 319 (2005).
- [12] A. Deandrea, G. Nardulli and A. Polosa, Phys. Rev. **D68**, 097501 (2003).
- [13] P. Colangelo and F. De Fazio, Phys. Lett. **B570**, 180 (2003).
- [14] S. Godfrey, Phys. Lett. **B568**, 254 (2003).
- [15] Fayyazuddin and Riazuddin, Phys. Rev. **D69**, 114008 (2004).
- [16] W. Wei, P.-Z. Huang and S.-L. Zhu, hep-ph/0510039.
- [17] T. Barnes, F.E. Close and H.J. Lipkin, Phys. Rev. **D68**, 054006 (2003).
- [18] A.P. Szczepaniak, Phys. Lett. **B567**, 23 (2003).
- [19] E. van Beveren and G. Rupp, Phys. Rev. Lett. **91**, 012003 (2003).
- [20] H.-Y. Cheng and W.-S. Hou, Phys. Lett. **B566**, 193 (2003).
- [21] K. Terasaki, Phys. Rev. **D68**, 011501(R) (2003).
- [22] L. Maiani, F. Piccinini, A.D. Polosa, V. Riquer, Phys. Rev. **D71**, 014028 (2005).
- [23] M.E. Bracco *et al.*, Phys. Lett. **B624**, 217 (2005).
- [24] H. Kim and Y. Oh, hep-ph/0508251.
- [25] T. Browder, S. Pakvasa and A.A. Petrov, Phys. Lett. **B578**, 365 (2004).
- [26] U. Dmitrasinovic, Phys. Rev. **D70**, 096011 (2004); Phys. Rev. Lett. **94**, 162002 (2005).
- [27] A. Hayashigaki and K. Terasaki, hep-ph/0410393.
- [28] R.L. Jaffe and F. Wilczek, Phys. Rev. Lett. **91**, 232003 (2003).
- [29] M.A. Shifman, A.I. and Vainshtein and V.I. Zakharov, Nucl. Phys. **B147**, 385 (1979).
- [30] L.J. Reinders, H. Rubinstein and S. Yazaky, Phys. Rep. **127**, 1 (1985).
- [31] B. L. Ioffe, Nucl. Phys. **B188**, 317 (1981); **B191**, 591(E) (1981).
- [32] S. Narison, Spectral Sum Rules, World Scientific Lecture Notes in Physics **26**.
- [33] M.E. Bracco, F.S. Navarra, M. Nielsen, Phys. Lett. **B454**, 346 (1999).
- [34] T.V. Brito, F.S. Navarra, M. Nielsen, M.E. Bracco, Phys. Lett. **B608**, 69 (2005).
- [35] M.E. Bracco *et al.*, Phys. Lett. **B521**, 1 (2001).
- [36] H.G. Dosch, E.M. Ferreira, F.S. Navarra and M. Nielsen, Phys. Rev. **D65**, 114002 (2002).
- [37] B. L. Ioffe and A.V. Smilga, Nucl. Phys. **B232**, 109 (1984).
- [38] M. Eidemüller *et al.*, Phys. Rev. **D72**, 034003 (2005).
- [39] J. Gasser, H. Leutwyler, Nucl. Phys. . **B250**, 465 (1985).
- [40] X. Jin, M. Nielsen, J. Pasupathy, Phys. Rev. **D51**, 3688 (1995); H. Forkel, M. Nielsen, Phys. Rev. **D55**, 1471 (1997), and references therein.
- [41] H.-C. Kim, M.M. Musakhanov, M. Siddikov, hep-ph/0508211.
- [42] I. Bediaga, M. Nielsen, Phys. Rev. **D68**, 036001 (2003).